


Agromet Advisory Service bulletin 14/17 Tuesday dt 04-04-17

(Issued jointly by IAAS Vellanikkara)





PALAKKAD

CURRENT SYNOPTIC SITUATION: .An upper air trough runs from comorin area to coastal Andhra Pradesh across Tamil Nadu and extends upto 0.9 km asl . An upper air cyclonic circulation lies over South Tamil Nadu and Neighbourhood between 1.5km and 2.1km a.s.l..

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 04.04.2017 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE: KERALA

PARAMETERS	ENSEMBLE FCST				
	DAY-1	DAY-2	DAY-3	DAY-4	DAY-
05/04		06/04	07/04	08/04	09/04
DISTRICT : PALAKKAD					
 Rainfall (mm)	1	0	0	0	
Max Temperature (deg C)	39	39	39	39	3:
Min Temperature (deg C)	26	26	26	26	2
Total cloud cover (octa)	4	4	6	6	
Max Relative Humidity (%)	85	85	85	85	8:
Min Relative Humidity (%)	45	45	45	45	4.
Wind speed (kmph)	006	006	006	004	00
Wind direction (deg)	050	050	050	050	05

Agro Meteorological Advisories:

Crop	Stage of the crop / Pest/Disease	Advisories
Paddy		Apply neem based pesticides (3 ml per one liter of water).
	Milky stage/Rice	Pesticides application should be done either before 9 am or
	bug	after 3 pm. Otherwise, the pesticide application will
		adversely affect the pollination.
Banana		Apply 2% neem emulsion in the starting stage. In case of
		severe attack apply 2.5 ml Chlorpyrifos in 1 liter of water
		on the pseudostem mixing with gum.
	Pseudo stem weevil	For destroying the weevil, Pseudo stem traps can be used.
		Split pseudostem traps (45 cm long) are made from the
		pseudostem pieces cut longitudinally in two halves, spread
		some amount of Beauveria in between them and a small
		piece of stone or pebble is placed between the stump and
		tie together & lay in the field. (Set 25 traps per acre).
Vegetables (Bitter gourd)		There is chance of mosaic in bitter gourd. Cut and destroy
	Mosaic	the infected plants. To control the vector of mosaic spray
		dimethoate (Roger) 1.5 ml /L.