


Advisory Service bulletin 15/17, Friday dt. 13-04-2017

(Issued jointly by IAAS Vellanikkara)

Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara E mail: kauagmet@yahoo.co.in, cohagmet@kau.in



PALAKKAD

CURRENT SYNOPTIC SITUATION: An upper air trough runs from north interior Karnataka to south Tamil Nadu extending upto 0.9km asl.

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 13.04.2017 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE: KERALA

PARAMETERS	ENSEMBLE FCST				
	DAY-1 14/04	DAY-2 15/04	DAY-3 16/04	DAY-4 17/04	DAY-5 18/04
ISTRICT : PALAKKAD					
ainfall (mm)	0	0	0	0	0
lax Temperature (deg C)	36	37	37	37	37
in Temperature (deg C)	27	27	27	27	26
otal cloud cover (octa)	6	6	6	5	5
ax Relative Humidity (%)	85	85	85	85	85
in Relative Humidity (%)	50	50	50	50	50
ind speed (kmph)	006	006	006	006	006
ind direction (deg)	270	270	270	270	270

Agro Meteorological Advisories:

Crop	Stage of the crop / Pest/Disease	Advisories		
Paddy	Milky stage/Rice bug	Apply neem based pesticides (3 ml per one liter of water). Pesticides application should be done either before 9 am or after 3 pm. Otherwise, the pesticide application will adversely affect the pollination.		
Coconut	Red palm weevil	There is a chance for red palm weevil in small coconut trees. As a control measure, mix 1 ml of imidachloprid in one litre of water and pour on its crown.		
Banana	-	Continue irrigation. Straw Mulching is recommended to conserve moisture. Destroy the side suckers.		
Vegetables (Amaranthus)	Leaf spot disease	Cut and remove the severely affected leaves. Apply 15 gm of Pseudomonas in cleared cow dung water after harvesting. Also apply Trichoderma mixed with cowdung in soil.		